

ON MODULAR INVERSES OF CYCLOTOMIC POLYNOMIALS AND THE MAGNITUDE OF THEIR COEFFICIENTS

CLÉMENT DUNAND

ABSTRACT. Let p and r be two primes and n, m be two distinct divisors of pr . Consider Φ_n and Φ_m , the n -th and m -th cyclotomic polynomials. In this paper, we present lower and upper bounds for the coefficients of the inverse of Φ_n modulo Φ_m .

1. INTRODUCTION

The magnitude of coefficients of polynomials derived from cyclotomic polynomials has attracted attention since the 19th century. If φ denotes the Euler totient function, the n -th cyclotomic polynomial Φ_n is a monic polynomial of degree $\varphi(n)$ whose roots are the primitive n -th roots of unity. In the following, we denote by a_i its coefficients.

Many results published so far deal with Φ_n . On one hand, we have asymptotic results which outline that these coefficients may have an exponential behavior for infinitely many n (see for instance Erdős [10] or Bateman [4]). On the other hand, there exists numerous studies for integers n composed of only few prime factors. In this way, Migotti [13] showed in 1883 that if n is composed of at most two primes p and r , the coefficients of Φ_{pr} can only be 0, -1 , 0 or 1. Later, around 1965, Beiter [5] and Carlitz [7] exhibit more precise criteria for these coefficients to be 0 or ± 1 . More recently, Lam and Leung [12] give in 1996 these coefficients in an explicit way.

The first example of a cyclotomic polynomial with a 2 among its coefficients is for $n = 105$. Yet, when n is the product of few primes, we can still find interesting bounds for the coefficients of Φ_n . It started in 1895 with Bang. For n product of three distinct primes, $p < q < r$, Bang [3] showed that $|a_i| \leq p - 1$. Later, in 1968, Beiter [5] gave a better bound when q or r equals ± 1 modulo p , that is $|a_i| \leq (p - 1)/2$. The conjecture that this bound could hold for all prime numbers p, q, r has been recently proved to be wrong by Gallot and Moree in [11]. Bachman [2] gave a better bound in 2003 for any p, q, r , $|a_i| \leq p - \lceil p/4 \rceil$. In 1968, Bloom [6] even gave a bound for $n = pqrs$, product of four primes with $p < q < r < s$: $|a_i| \leq p(p - 1)(pq - 1)$.

This paper deals with modular inverses of cyclotomic polynomials. If Φ_m and Φ_n are coprime, then Φ_m is invertible modulo Φ_n and, following the example of Φ_n , we may ask whether the coefficients of $\Phi_m^{-1} \bmod \Phi_n$ are of special form. Especially, we noticed that the magnitude of these coefficients is very particular when n is composed of few prime factors, and we thoroughly prove lower and upper bounds for them when m and n are two distinct

Date: July 31, 2009.

divisors of pr , the product of two primes. In the case of the product of three primes pqr , the particular structure of Φ_{pqr} may also yield interesting results, but is out of the scope of this work.

Our main motivation is the computation of the isomorphism between the multiplicative group of a finite field $\mathbb{F}_{q^N}$ and the product of some of its subgroups, since $q^N - 1 = \prod_{d|N} \Phi_d(q)$. Such calculations typically occur in torus-based cryptographic schemes, as developed by Silverberg and Rubin [14], and the present bounds lead improvements in the running times of algorithms in this field (see [8, ?]). But, these inverses show up in many other contexts, since they are inverses of cyclotomic units in cyclotomic fields.

Section 2 is dedicated to the resultant of Φ_m and Φ_n . To that end, we show the following lemma.

Lemma 1. *For all integers $m > n \geq 1$,*

$$\text{Res}(\Phi_m, \Phi_n) \neq 1 \Leftrightarrow m = np^\alpha \quad \text{with } p \text{ prime and } \alpha \geq 1.$$

This is a consequence of a result by Apostol [1] about the resultant of cyclotomic polynomials. We suspect that it is already known since it helps proving the equivalence of two definitions of an algebraic torus (see [14] again for further details) but we did not find any explicit proof of it in the literature. As a result, we obtain a condition for the coprimality of $\Phi_m(q)$ and $\Phi_n(q)$, whatever the integer q is.

In the case of two coprime cyclotomic polynomials, we can consider the inverse of Φ_m modulo Φ_n . In Section 3, we make an exhaustive study when n and m are divisors of the product of two primes.

Theorem 1. *For all p and r distinct prime numbers,*

- (i) $\Phi_p^{-1} \bmod \Phi_1 = 1/p$ and $\Phi_1^{-1} \bmod \Phi_p = (-1/p)(X^{p-2} + 2X^{p-3} + \dots + p - 1)$.
- (ii) $\Phi_{pr}^{-1} \bmod \Phi_1 = 1$ and $\Phi_1^{-1} \bmod \Phi_{pr} = \sum_{i=0}^{\varphi(pr)-1} v_i X^i$ with $v_i \in \{-1, 0, 1\}$.
- (iii) $\Phi_{pr}^{-1} \bmod \Phi_p = \frac{1}{r} \sum_{i=0}^d X^i$ with $d \equiv r-1 \bmod p$ and $\Phi_p^{-1} \bmod \Phi_{pr} = \frac{1}{r} \sum_{i=0}^{\varphi(pr)-1} v_i X^i$ with $v_i < r$.
- (iv) $\Phi_p^{-1} \bmod \Phi_r = \sum_{i=0}^{\varphi(r)-1} v_i X^i$ with $v_i \in \{0, -1, +1\}$.

Notations. In this paper, $\mathcal{P}$ denotes the set of all prime numbers and (m, n) is the greatest common divisor of m and n . We also recall the following result about Möbius μ function,

$$\forall n > 1, \quad \sum_{d|n} \mu(d) = 0. \quad (1)$$

2. COPRIMALITY

In this section, we consider the resultant of two cyclotomic polynomials Φ_m and Φ_n . In order to prove Lemma 1, we start from Theorem 2, due to Apostol [1].

Theorem 2 (Apostol, [1]). *Let $m > 1$, then*

$$\text{Res}(\Phi_1, \Phi_m) = \begin{cases} p & \text{if } m = p^a, p \text{ prime, } a \geq 1, \\ 1 & \text{otherwise.} \end{cases}$$

Besides, let $m > n > 1$, then

$$\text{Res}(\Phi_m, \Phi_n) = \prod_{\substack{d|n \\ p \in \mathcal{P} \text{ such that } \frac{m}{(m,d)} = p^a}} p^{\mu(n/d) \frac{\varphi(m)}{\varphi(p^a)}} \quad (2)$$

where μ is the Möbius function and φ the d'Euler totient function. This product is performed over the divisors d of n such that $m/(m, d)$ is a prime power p^a .

We can now prove Lemma 1.

Proof. For $n = 1$, we have $\text{Res}(\Phi_m, \Phi_1) = (-1)^{\varphi(m)} \text{Res}(\Phi_1, \Phi_m)$ and $\varphi(m)$ is even as soon as $m > 2$. So the resultant equals 1 if and only if m is not a prime power.

Now, let us consider $m > n > 1$. This time we are going to use Eq.(2).

Sufficiency. If $m = np^\alpha$, we can consider the powers of p showing up in the product and show that it does not equal 1.

For d dividing n , we have $m/(m, d) = (n/d)p^\alpha$ because $n|m$. So this is a power of p , only if $n/d = p^\varepsilon$. But $\mu(n/d) = 0$ as soon as $\varepsilon > 1$. So the only non trivial terms will correspond to the cases $\varepsilon = 1$ and $\varepsilon = 0$.

- For $\varepsilon = 0$, we have $d = n$, so $m/(m, d) = p^\alpha$ and $a = \alpha$, which implies

$$p^{\mu(\frac{n}{d}) \frac{\varphi(m)}{\varphi(p^\alpha)}} = p^{\frac{\varphi(m)}{\varphi(p^\alpha)}}.$$

- For $\varepsilon = 1$, we have $d = n/p$, so $m/(m, d) = p^{\alpha+1}$ and $a = \alpha + 1$, which implies

$$p^{\mu(\frac{n}{d}) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} = p^{-\frac{\varphi(m)}{\varphi(p^{\alpha+1})}}.$$

The contribution in terms of powers of p is $p^{\varphi(m) \left(\frac{1}{\varphi(p^\alpha)} - \frac{1}{\varphi(p^{\alpha+1})} \right)} > 1$.

Necessity. We want to isolate the common factor of m and n , if they have one. That is to say, we write $m = wM$ and $n = wN$ with $(M, N) = 1$.

Since the resultant is not 1, we have at least one non trivial term $p^{\mu(n/d) \frac{\varphi(m)}{\varphi(p^a)}}$ in the product for some d such that $m/(m, d) = p^a$.

Since $d|wN$, we can write $d = d_1 d_2$ with $d_1|w$ and $d_2|N$. Let us suppose d_1 maximal for this property, i.e. $(w/d_1, d_2) = 1$. We call $w' = w/d_1$. Then

$$\frac{m}{(m, d)} = \frac{wM}{(wM, d_1 d_2)} = \frac{wM}{d_1 (w' M, d_2)} = \frac{wM}{d_1 (w', d_2)} = \frac{w}{(w, d)} M.$$

So $(w/(w, d))M = p^a$, which implies that M is a power of p , say $M = p^\alpha$. But $w/(w, d)$ is also a power of p . Let p^r be the greatest power of p in w , so that $w = p^r s$ with $p \nmid s$. Thus the powers involved in the product are $\mu(n/d) = \mu(p^r s N/d)$.

We know that it will be zero as soon as p has power at least 2 in μ . So the contribution to the product will be non trivial only if $d = p^{r-\varepsilon}\delta$ with $\varepsilon = 0$ or 1 if $r \geq 1$ and $\delta \mid sN$. More precisely $\delta = s'd_2$ with $d_2 \mid N$ and $s' \mid s$. Then $(w, d) = p^{r-\varepsilon}(s/s')$ and we can even assert that $s = s'$ because $w/(w, d)$ must be a power of p and $p \nmid s$.

Finally,

$$w = p^r s, \quad d = p^{r-\varepsilon} s d_2 \text{ with } d_2 \mid N \text{ and } \varepsilon = 0 \text{ or } 1 \text{ if } r \geq 1.$$

Thus $w/(w, d) = p^{a-\alpha} = p^\varepsilon$. Now we can give the contribution to the product.

If $\varepsilon = 0$, then $d = w d_2$, $a = \alpha$ and the product is

$$\prod_{d \mid n} p^{\mu(n/d) \frac{\varphi(m)}{\varphi(p^\alpha)}} = \prod_{d_2 \mid N} p^{\mu(N/d_2) \frac{\varphi(m)}{\varphi(p^\alpha)}} = \left(p^{\frac{\varphi(m)}{\varphi(p^\alpha)}} \right)^{\sum_{d_2 \mid N} \mu\left(\frac{N}{d_2}\right)} = \left(p^{\frac{\varphi(m)}{\varphi(p^\alpha)}} \right)^{\sum_{d_2 \mid N} \mu(d_2)}$$

If $\varepsilon = 1$, then $d = w d_2 / p$, $a = \alpha + 1$ and the product is

$$\begin{aligned} \prod_{d \mid n} p^{\mu(n/d) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} &= \prod_{d_2 \mid N} p^{\mu\left(\frac{pN}{d_2}\right) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} = \prod_{d_2 \mid N} p^{\mu(p) \mu\left(\frac{N}{d_2}\right) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} \quad \text{since } (p, N) = 1, \\ &= \left(p^{\mu(p) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} \right)^{\sum_{d_2 \mid N} \mu\left(\frac{N}{d_2}\right)} = \left(p^{\mu(p) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} \right)^{\sum_{d_2 \mid N} \mu(d_2)}. \end{aligned}$$

Thus,

$$\left(p^{\frac{\varphi(m)}{\varphi(p^\alpha)}} p^{\mu(p) \frac{\varphi(m)}{\varphi(p^{\alpha+1})}} \right)^{\sum_{d_2 \mid N} \mu(d_2)} \text{ divides } \text{Res}(\Phi_m, \Phi_n).$$

Since this result should be greater than 1, necessarily $\sum_{d_2 \mid N} \mu(d_2) \neq 0$ which is impossible unless $N = 1$ according to Eq. (1), and thus $n = w$ and $m = np^\alpha$. $\square$

Now it is easy to show the following condition of coprimality.

Corollary 1. *For any integer q and $m > n \geq 1$ integers, $\Phi_m(q)$ and $\Phi_n(q)$ are coprime if m does not divide n .*

Proof. If m does not divide n , we know from Lemma 1 that $\text{Res}(\Phi_m, \Phi_n) = 1$, which is true in $\mathbb{Z}$ but also in $\mathbb{Z}/\ell\mathbb{Z}$ for any $\ell \in \mathbb{Z}$ since 1 is unchanged. Now suppose that $\Phi_m(q)$ and $\Phi_n(q)$ have a common factor, say ℓ . Then Φ_m and Φ_n have a common root, q , in $\mathbb{Z}/\ell\mathbb{Z}$ and consequently their resultant is zero, which is false. $\square$

3. INVERSION OF $\Phi_m \bmod \Phi_n$

Consider m and n such that the cyclotomic polynomials Φ_m and Φ_n are coprime. Then Φ_m is invertible modulo Φ_n and it is a natural question to try to compute Φ_m^{-1} modulo Φ_n and more precisely we would like to know the magnitude of its coefficients.

Since Φ_m and Φ_n are coprime we can write the Bézout relation

$$\Phi_m U + \Phi_n V = 1.$$

Our goal is to find $U = \Phi_m^{-1} \bmod \Phi_n$ or at least quantify the magnitude of its coefficients.

In this section we are going to prove the four assertions of Theorem 1 in turn. We recall that $\Phi_n(1) = p$ if $n = p^\alpha$ is a prime power; else $\Phi_n(1) = 0$ for $n > 1$.

3.1. Case $m = p$ and $n = 1$. The cyclotomic polynomials Φ_p and Φ_1 are both easy to write and it is not difficult to obtain explicit expressions for their inverses.

Proposition 1. *For all prime number p ,*

- $\Phi_p^{-1} \bmod \Phi_1 = 1/p$,
- $\Phi_1^{-1} \bmod \Phi_p = -\frac{1}{p}(X^{p-2} + 2X^{p-3} + \dots + p - 1)$.

Proof. We simply check that the Bézout relation between Φ_p and Φ_1 is valid.

$$\begin{aligned} -\Phi_1(X)(X^{p-2} + 2X^{p-3} + \dots + p - 1) + \Phi_p(X) &= (X - 1) \sum_{k=0}^{p-2} (k + 1 - p)X^k + \Phi_p(X) \\ &= \sum_{k=1}^{p-1} (k - p)X^k - \sum_{k=0}^{p-2} (k + 1 - p)X^k + \sum_{k=0}^{p-1} X^k = p. \end{aligned}$$

□

3.2. Case $m = pr$ and $n = 1$. The explicit expression of Φ_{pr} is less convenient than that of Φ_p , but we still have useful information thanks to Lam and Leung [12].

Proposition 2. *For all p and r distinct prime numbers,*

- $\Phi_{pr}^{-1} \bmod \Phi_1 = 1$.
- $\Phi_1^{-1} \bmod \Phi_{pr} = \sum_{i=0}^{(p-1)(r-1)-1} v_i X^i$ with $v_i \in \{-1, 0, 1\}$.

Proof. We are first looking for U in the Bézout relation $\Phi_{pr}U + \Phi_1V = 1$ and we know that it has degree 0. So a simple evaluation of this relation at 1 gives $U(1) = 1$, so $\Phi_{pr}^{-1} \bmod \Phi_1 = 1$ because $\Phi_{pr}(1) = 1$.

Now V is characterized by $(X - 1)V(X) = 1 - \Phi_{pr}(X)$. Let $V(X) = \sum_{i=0}^d v_i X^i$ and $\Phi_{pr}(X) = \sum_{i=0}^d a_i X^i$ with $d = (p - 1)(r - 1) - 1$. Then we can write the equation as a linear system,

$$\begin{bmatrix} -1 & 0 & \dots & 0 \\ 1 & -1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 1 & -1 \end{bmatrix} \begin{bmatrix} v_0 \\ v_1 \\ \vdots \\ v_d \end{bmatrix} = \begin{bmatrix} 1 - a_0 \\ -a_1 \\ \vdots \\ -a_d \end{bmatrix}, \text{ equivalent to } \begin{bmatrix} v_0 \\ v_1 \\ \vdots \\ v_d \end{bmatrix} = \begin{bmatrix} 1 & 0 & \ddots & 0 \\ 1 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 1 & \dots & 1 & 1 \end{bmatrix} \begin{bmatrix} a_0 - 1 \\ a_1 \\ \vdots \\ ta_d \end{bmatrix}.$$

Note that we know from [12] that $a_0 = 1$ and for all i , $a_i \in \{0, \pm 1\}$. Moreover the signs (+1 or -1) are alternating. So each v_i is necessarily 0 or ± 1 . □

Note that a similar technique could allow us to compute Φ_n^{-1} modulo Φ_1 for any n since it is simply $\Phi_n(1)$ which we explicitly know.

3.3. Case $m = pr$ and $n = p$. This time we will need the explicit expression of Φ_{pr} .

Proposition 3. *For all p and r distinct prime numbers,*

$$\Phi_{pr}^{-1} \bmod \Phi_p = \frac{1}{r} \sum_{i=0}^d X^i \text{ with } d \equiv r-1 \bmod p.$$

Proof. Let us directly show that $\frac{1}{r} \left(\sum_{i=0}^d X^i \right) \Phi_{pr} \equiv 1 \bmod \Phi_p$. For this purpose, we need to use the expression of Φ_{pr} given in [12]. Let s and t be two integers such that $(p-1)(r-1) = \varphi(pr) = sp + tq$. Then,

$$\Phi_{pr}(X) = \left(\sum_{i=0}^s X^{ip} \right) \left(\sum_{j=0}^t X^{jr} \right) - \left(\sum_{i=s+1}^{r-1} X^{ip} \right) \left(\sum_{j=t+1}^{p-1} X^{jr} \right) X^{-pr}.$$

Thus,

$$\begin{aligned} \Phi_{pr} \sum_{i=0}^d X^i \bmod \Phi_p &= \left((s+1) \sum_{j=0}^t X^{jr} - (r-1-s) \sum_{j=t+1}^{p-1} X^{jr} \right) \sum_{i=0}^d X^i \\ &= \left((s+1) \sum_{j=0}^{p-1} X^{jr} - r \sum_{j=t+1}^{p-1} X^{jr} \right) \sum_{i=0}^d X^i \end{aligned}$$

But $\Phi_p(X^r) = \sum_{j=0}^{p-1} X^{jr}$, and then

$$\begin{aligned} \Phi_{pr} \sum_{i=0}^d X^i \bmod \Phi_p &= -r \sum_{j=t+1}^{p-1} X^{jr} \sum_{i=0}^d X^i \\ &= r \frac{X^{(t+1)r} - 1}{X^r - 1} \frac{X^{d+1} - 1}{X - 1} \end{aligned}$$

An explicit computation shows that $(t+1)r = 1 + pr - p(s+1)$. So $X^{(t+1)r} \equiv X \bmod \Phi_p$. Besides $d+1 \equiv r \bmod p$, so $X^{d+1} \equiv X^r \bmod \Phi_p$, which leads to the result. There only remains r in the computed product. $\square$

3.4. Case $m = p$ and $n = pr$. Then the proof consists in examining the Euclidean division step by step.

Proposition 4. *For all p and r distinct prime numbers,*

$$\Phi_p^{-1} \bmod \Phi_{pr} = \frac{1}{r} \sum_i v_i X^i \text{ with } v_i < r.$$

Proof. We are looking for V in the Bézout relation $\Phi_{pr}U + \Phi_pV = 1$, where $U = \frac{1}{r} \sum_{i=0}^d X^i$ from Proposition 3. So, all we have to do is divide $1 - \Phi_{pr}U$ by Φ_p .

First, note that $1 - \Phi_{pr}U$ has only coefficients ± 1 . Indeed we know that the coefficients of Φ_{pr} are alternating $+1$'s and -1 's among other 0 's. So, if we write their explicit product, we

obtain a polynomial with simple coefficients (only 0 or ± 1) thanks to the Cauchy product. In fact, we divide a polynomial with coefficients in $\{0, \pm 1\}$ by Φ_p .

If we simply examine the Euclidean division step by step, we can show by recurrence that the range of possible coefficients for the quotient increases by one at each step, and always contains 0. $\square$

3.5. Case $m = p$ and $n = r$ with p and r two distinct primes. Before we give a proof for the last assertion in Theorem 1, we need to work on the general problem. The idea is to evaluate our Bézout relation at the roots of Φ_n and to interpolate U from the values found at these points.

The roots of Φ_n are the primitive n -th roots of 1. If we call ξ one of them and I_n the set of integers prime with n : $I_n = \{1 \leq j \leq n-1, (j, n) = 1\}$, they are $\{\xi^j, j \in I_n\}$.

The evaluation of our Bézout relation in these points gives

$$\forall j \in I_n, \quad \Phi_m(\xi^j)U(\xi^j) = 1.$$

If we note $U = \sum_{i=1}^{\varphi(n)} u_i X^{i-1}$ then the equation can be written

$$\forall j \in I_n, \quad \sum_{i=1}^{\varphi(n)} u_i (\xi^j)^{i-1} = (\Phi_m(\xi^j))^{-1}.$$

Proposition 5. *For all distinct prime numbers p and r ,*

$$\Phi_p^{-1} \bmod \Phi_r = \sum_{i=1}^{\varphi(r)} u_i X^{i-1} \text{ with } u_i \in \{-1, 0, +1\}.$$

Proof. We show in appendix A (Lemma 3) that

$$u_i = \sum_{j=1}^{r-1} (-1)^{i+j} \sigma_{r-1-i}(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1}) \frac{V(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1})}{V(\xi, \dots, \xi^{r-1})} \Phi_p(\xi^j)^{-1}.$$

The primality of p and r yields very convenient simplifications: $I_r = \{1, \dots, r-1\}$ and $\varphi(r) = r-1$, but also $\Phi_p(X) = 1 + X + \dots + X^{p-1}$ and so $\Phi_p(1) = p$. Similarly $\Phi_r(1) = r$. Let us first calculate the quotient of Vandermonde determinants.

$$\begin{aligned} \frac{V(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1})}{V(\xi, \dots, \xi^{r-1})} &= \prod_{\substack{1 \leq k \leq r-1 \\ k < j}} (\xi^j - \xi^k)^{-1} \prod_{\substack{1 \leq k \leq r-1 \\ k > j}} (\xi^k - \xi^j)^{-1} \\ &= \frac{(-1)^{r-1-j}}{(\xi^j)^{r-2}} \prod_{\substack{1 \leq k \leq r-1 \\ k \neq j}} (1 - \xi^{k-j})^{-1} \\ &= \frac{(-1)^{r-1-j}}{(\xi^j)^{-2} (1 - \xi^{-j})^{-1}} \prod_{\substack{0 \leq k \leq r-1 \\ k \neq j}} (1 - \xi^{k-j})^{-1} \end{aligned}$$

But the remaining product is $\Phi_r(1)^{-1} = 1/r$. Thus,

$$\frac{V(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1})}{V(\xi, \dots, \xi^{r-1})} = \frac{(-1)^j}{r} \xi^j (\xi^j - 1).$$

We can write $\Phi_p(X)$ as a geometric sum. So $\Phi_p(\xi^j)^{-1} = (1 - \xi^j)/(1 - \xi^{jp})$. Finally we can give an explicit expression of $\sigma_{r-1-i}(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1})$. Indeed,

$$\prod_{\substack{1 \leq k \leq r-1 \\ k \neq j}} (X - \xi^k) = \sum_{i=1}^{r-1} (-1)^{r-1-i} \sigma_{r-1-i}(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1}) X^{i-1}.$$

So we simply need to find the coefficients of $\prod_{\substack{1 \leq k \leq r-1 \\ k \neq j}} (X - \xi^k) = \frac{\Phi_r(X)}{(X - \xi^j)}$.

The following calculation shows that $\sigma_{r-1-i}(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1}) = (-1)^i \sum_{k=0}^{i-1} (\xi^j)^k$.

$$\begin{aligned} (X - \xi^j) \sum_{i=1}^{r-1} \left(\sum_{k=0}^{i-1} (\xi^j)^k \right) X^{r-1-i} &= \sum_{i=1}^{r-1} \left(\sum_{k=0}^{i-1} (\xi^j)^k \right) X^{r-i} - \sum_{i=1}^{r-1} \left(\sum_{k=0}^{i-1} (\xi^j)^{k+1} \right) X^{r-1-i} \\ &= \sum_{i=0}^{r-2} \left(\sum_{k=0}^i (\xi^j)^k \right) X^{r-1-i} - \sum_{i=1}^{r-1} \left(\sum_{k=1}^i (\xi^j)^k \right) X^{r-1-i} \\ &= X^{r-1} + \underbrace{\sum_{i=1}^{r-2} \left(\sum_{k=0}^i (\xi^j)^k - \sum_{k=1}^i (\xi^j)^k \right)}_{=1} X^{r-1-i} - \sum_{k=1}^{r-1} (\xi^j)^k \\ &= \Phi_r(X). \end{aligned}$$

And this expression can be seen as a geometric sum again.

$$\sigma_{r-1-i}(\xi, \dots, \hat{\xi}^j, \dots, \xi^{r-1}) = (-1)^i \frac{1 - \xi^{-ij}}{1 - \xi^j}.$$

Now we can simplify our expression for u_i :

$$\begin{aligned} u_i &= \frac{1}{r} \sum_{j=1}^{r-1} \frac{(-1)^{i+j} (-1)^i (-1)^j (1 - \xi^{-ij}) \xi^j (\xi^j - 1) (1 - \xi^j)}{(1 - \xi^j) (1 - \xi^{jp})} \\ &= \frac{1}{r} \sum_{j=1}^{r-1} \frac{\xi^j (1 - \xi^{-ij}) (\xi^j - 1)}{(1 - \xi^{jp})} \end{aligned}$$

We can improve this expression using the following relation:

$$\frac{1}{1 - \xi^{jp}} = \frac{1}{r} \left(\xi^{jp(r-2)} + 2\xi^{jp(r-3)} + \dots + (r-1) \right).$$

Indeed,

$$\begin{aligned}
(1 - \xi^{jp}) \sum_{k=1}^{r-1} k \xi^{jp(r-1-k)} &= \sum_{k=1}^{r-1} k \xi^{jp(r-1-k)} - \sum_{k=1}^{r-1} k \xi^{jp(r-k)} \\
&= \sum_{k=2}^r (k-1) \xi^{jp(r-k)} - \sum_{k=1}^{r-1} k \xi^{jp(r-k)} \\
&= (r-1) - \underbrace{\sum_{k=1}^{r-2} (\xi^{jp})^k}_{=-1 - \xi^{jp(r-1)} \text{ if } jp \not\equiv 0 \pmod r} - \xi^{jp(r-1)} \\
&= r \quad \text{since } jp \not\equiv 0 \pmod r \text{ (} p \text{ prime and } 1 \leq j \leq r-1 \text{)}.
\end{aligned}$$

So the final expression for u_i is

$$\begin{aligned}
u_i &= \frac{-1}{r^2} \sum_{j=1}^{r-1} \xi^j (1 - \xi^{-ij}) (1 - \xi^j) \sum_{k=1}^{r-1} k \xi^{jp(r-1-k)} \\
&= \frac{-1}{r^2} \sum_{j=1}^{r-1} \xi^j (1 - \xi^{-ij}) (1 - \xi^j) \sum_{k=0}^{r-2} (r-k-1) \xi^{jp k} \\
&= \frac{-1}{r^2} \sum_{k=0}^{r-2} (r-k-1) \sum_{j=1}^{r-1} (\xi^{j(pk+1)} - \xi^{j(pk+2)} - \xi^{j(pk-i+1)} + \xi^{j(pk-i+2)}) \\
&= \frac{-1}{r^2} \sum_{k=0}^{r-2} (r-k-1) \underbrace{(S_1(k) - S_2(k) - S_3(k) + S_4(k))}_{S(k)}.
\end{aligned}$$

The sums $S_l = \sum_{j=1}^{r-1} (\xi^A)^{jl}$ are actually sums of all the powers of a r -th root of 1, except $(\xi^A)^0 = 1$. So if ξ^A is a prime root of 1, the sum simply equals -1. And if ξ^A is not a prime root of 1, the only possibility is $\xi^A = 1$ (*i.e.* $A \equiv 0 \pmod r$) and in this case the sum equals $r-1$.

So u_i depends on the value of the powers $pk+1$, $pk+2$, $pk-i+1$ and $pk-i+2$ modulo q .

Most of time the four sums involved will all be equal to 1 and thus $S_1 + S_2 + S_3 + S_4 = 0$. But there can be up to four values of k for which one of the four sums will not be equal to 1 but to $r-1$.

- If there exists $0 \leq k_1 \leq r-2$ such that $1 + pk_1 \equiv 0 \pmod r$ then $S_1(k_1) = r-1$.
- If there exists $0 \leq k_2 \leq r-2$ such that $2 + pk_2 \equiv 0 \pmod r$ then $S_2(k_2) = r-1$.
- If there exists $0 \leq k_3 \leq r-2$ such that $1 - i + pk_3 \equiv 0 \pmod r$ then $S_3(k_3) = r-1$.
- If there exists $0 \leq k_4 \leq r-2$ such that $2 - i + pk_4 \equiv 0 \pmod r$ then $S_4(k_4) = r-1$.

The most important argument now is the following : in each sum S_l all the powers of ξ appear, except the power involving $k = r-1$. In other words, for S_1 for instance, all the

sums $S_1(k)$ appear, for $0 \leq k \leq r-2$. Besides the $\{1 + pk \bmod r, 0 \leq k \leq r-1\}$ take all values $\{0, \dots, r-1\}$ because p and r are coprime. So either there will exist $0 \leq k_1 \leq r-2$ such that $1 + pk_1 \equiv 0 \bmod r$ or necessarily $1 + p(r-1) \equiv 0$.

Let us first prove that at least three k 's among k_1, k_2, k_3 and k_4 exist. If there does not exist a $k_j, j = 1, 2, 3$ or 4 , then $p(r-1) + l \equiv 0[r]$ with $l \in \{1, 2, 1-i, 2-i\}$ respectively. So if two different k 's do not exist, we will have this relation for l and l' among $\{1, 2, 1-i, 2-i\}$. So $l \equiv l' \bmod r$, which is possible only if $i = 1$ or $i = r-1$.

So we have proved that at least three of the four k_j 's exist except for $i = 1$ and $r-1$. In the last two cases, two k_j 's at least still exist since $p(r-1) + l \equiv 0[r]$ cannot be simultaneously true for more than two different $l \in \{1, 2, i-1, i-2\}$.

Let us first consider the particular case $i = r-1$. Then $S = S_1 - 2S_2 + S_4$.

The possible contributions to the sum S for $k = k_1, k_2$ or k_4 are $S(k_1) = (r-1) - 2(-1) + (-1) = r$ and similarly $S(k_2) = -2r$ and $S(k_4) = r$.

We know that at least 2 of these k 's exist. So we have four different possibilities.

- If there exist k_1, k_2 and k_4 then we have a system of three equations

$$\begin{cases} 1 + pk_1 & \equiv 0[r] & (1a) \\ 2 + pk_2 & \equiv 0[r] & (1b) \\ 3 + pk_4 & \equiv 0[r] & (1c) \end{cases}.$$

So r divides $p(2k_2 - k_1 - k_4)$ and thus $r | (2k_2 - k_1 - k_4)$. Since each $|k_i|$ does not exceed $r-1$, $2k_2 - k_1 - k_4 = 0$ or $\pm r$.

So $r^2 u_1 = r(r - k_1 - 1) - 2r(r - k_2 - 1) + r(r - k_4 - 1) = r(2k_2 - k_1 - k_4)$. Thus $u_1 = 0$ or ± 1 .

- If k_1 does not exist then equation (1a) is replaced by $1 + p(r-1) \equiv 0 \bmod r$. So $p \equiv 1 \bmod r$. Then solving equations (1b) and (1c) with this particular p gives $k_2 = r-2$ and $k_4 = r-3$.

So $r^2 u_1 = [(r - k_2 - 1)(-2r) + (r - k_4 - 1)r] = -2r^2 + 2r^2$. Thus $u_1 = 0$.

- If k_2 does not exist then equation (1b) is replaced by $2 + p(r-1) \equiv 0 \bmod r$. So $p \equiv 2 \bmod r$. Then solving equations (1a) and (1c) with this particular p gives $2k_1 = r-1$ and $2k_4 = r-3$.

So $2r^2 u_1 = [(2r - 2k_1 - 2)r + (2r - 2k_4 - 2)r] = 2r^2$. Thus $u_1 = 1$.

- If k_4 does not exist then equation (1c) is replaced by $3 + p(r-1) \equiv 0 \bmod r$. So $p \equiv 3 \bmod r$. Then solving equations (1a) and (1b) with this particular p gives

$$\begin{cases} 3k_1 \equiv -1 \bmod r \\ 3k_2 \equiv -2 \bmod r \end{cases} \Rightarrow \begin{cases} 3k_1 = r-1 \\ 3k_2 = 2r-2 \end{cases} \quad \text{or} \quad \begin{cases} 3k_1 = 2r-1 \\ 3k_2 = r-2 \end{cases}$$

since $r-1$ and $r-2$ cannot be both divisible by 3.

$$\begin{aligned} \text{So } 3r^2 u_1 &= [(3r - 3k_1 - 3)r + (3r - 3k_2 - 3)(-2r)] \\ &= (2r - 2)r + (r - 1)(-2r) \text{ or } (r - 2)r + (2r - 1)(-2r). \end{aligned}$$

Thus $u_1 = 0$ or -1 .

Now let us consider the particular case $i = 1$. It is very similar to the previous case. Then $S = 2S_1 - S_2 - S_3$, where $S_3 = \sum_{j=1}^{r-1} \xi^{j(pk)}$. So the powers involved are pk , $pk + 1$ and $pk + 2$.

The possible contributions to the sum S for $k = k_1, k_2$ or k_3 are $S(k_1) = 2r$, $S(k_2) = -r$ and $S(k_3) = -r$. Note that $k_3 = 0$ since it is such that $pk_3 \equiv 0 \pmod{r}$.

We know that at least 2 of these k 's exist. So we have three different possibilities.

- If there exist k_1, k_2 and k_3 then we have the system of three equations

$$\begin{cases} 1 + pk_1 & \equiv 0[r] & (2a) \\ 2 + pk_2 & \equiv 0[r] & (2b) \\ pk_3 & \equiv 0[r] & (2c) \end{cases}.$$

So r divides $p(2k_2 - k_1 - k_4)$ and thus $r | (2k_2 - k_1 - k_4)$. Since each $|k_i|$ does not exceed $r - 1$, $k_2 - 2k_1 = 0$ or $\pm r$. So $r^2 u_{r-1} = 2r(r - k_1 - 1) - r(r - k_2 - 1) - r(r - k_3 - 1) = r(k_2 - 2k_1)$. Thus $u_{r-1} = 0$ or ± 1 .

- If k_1 does not exist then equation (2a) is replaced by $1 + p(r - 1) \equiv 0 \pmod{r}$. So $p \equiv 1 \pmod{r}$. Then solving equations (2b) and (2c) with this particular p gives $k_2 = r - 2$ and $k_3 = 0$.

So $r^2 u_{r-1} = [(r - k_2 - 1)(-r) + (r - k_3 - 1)(-r)] = -r - r(r - 1)$. Thus $u_{r-1} = -1$.

- If k_2 does not exist then equation (2b) is replaced by $2 + p(r - 1) \equiv 0 \pmod{r}$. So $p \equiv 2 \pmod{r}$. Then solving equations (2a) and (2c) with this particular p gives $2k_1 = r - 1$ and $2k_3 = 0$.

So $2r^2 u_{r-1} = [(2r - 2k_1 - 2)(2r) + (2r - 2)(-r)] = 0$. Thus $u_{r-1} = 0$.

Finally consider the general case $1 < i < r - 1$. The potential nonzero contributions to u_i are $S(k_1) = (r - 1) - 2(-1) + (-1) = r$ and similarly $S(k_2) = -r$, $S(k_3) = -r$ and $S(k_4) = r$ when respectively k_1, k_2, k_3 and k_4 exist.

As we have previously seen, at most one k does not exist. So we have five different cases.

- If k_1, k_2, k_3 and k_4 exist then we have the system of four equations

$$\begin{cases} 1 + pk_1 & \equiv 0[r] & (3a) \\ 2 + pk_2 & \equiv 0[r] & (3b) \\ 1 - i + pk_3 & \equiv 0[r] & (3c) \\ 2 - i + pk_4 & \equiv 0[r] & (3d) \end{cases}.$$

Since p and r are coprime, $r | (k_2 + k_3 - k_1 - k_4)$. But similarly to a previous case no $|k|$ exceeds $r - 1$. So finally $k_2 + k_3 - k_1 - k_4 = 0$ or $\pm r$.

So $r^2 u_i = r(k_2 + k_3 - k_1 - k_4) \Rightarrow u_i = 0$ or ± 1 .

- If k_1 does not exist then equation (3a) is replaced by $1 + p(r - 1) \equiv 0 \pmod{r}$. So $p \equiv 1 \pmod{r}$. Then solving equations (3b), (3c) and (3d) with this particular p gives $k_2 = r - 2$, $k_3 = i - 1$ and $k_4 = i - 2$.

So $r^2 u_1 = (r - k_2 - 1)(-r) + (r - k_3 - 1)(-r) + (r - k_4 - 1)r = 0$. Thus $u_i = 0$.

- If k_2 does not exist then equation (3b) is replaced by $2 + p(r - 1) \equiv 0 \pmod{r}$. So $p \equiv 2 \pmod{r}$. Then solving equations (3a), (3c) and (3d) with this particular p

gives

$$\begin{cases} 2k_1 \equiv -1 \pmod{r} \\ 2k_3 \equiv i-1 \pmod{r} \\ 2k_4 \equiv i-2 \pmod{r} \end{cases} \Rightarrow \begin{cases} 2k_1 = r-1 \\ 2k_3 = i-1 \\ 2k_4 = r+i-2 \end{cases} \quad \text{or} \quad \begin{cases} 2k_1 = r-1 \\ 2k_3 = r+i-1 \\ 2k_4 = i-2 \end{cases}$$

since $i-1$ and $i-2$ cannot be both even.

So $2r^2u_1 = (2r - 2k_1 - 2)r + (2r - 2k_3 - 2)(-r) + (2r - 2k_4 - 2)r = 0$ or $2r^2$.

Thus $u_i = 0$ or 1 .

- If k_3 does not exist then equation (3c) is replaced by $1 - i + p(r-1) \equiv 0 \pmod{r}$. So $p \equiv 1 - i \pmod{r}$. Then combining equations (3a), (3b) and (3d) with this particular p shows that $(1-i)(k_4 + k_1 - k_2 + 1) \equiv 0 \pmod{r}$. Since r and $1-i$ are coprime, $r|(k_4 + k_1 - k_2 + 1)$, thus $k_4 + k_1 - k_2 + 1 = 0$ or r .
So $(1-i)r^2u_i = (1-i)r^2$ or 0 . Thus $u_i = 1$ or 0 .
- If k_4 does not exist then equation (3d) is replaced by $2 - i + p(r-1) \equiv 0 \pmod{r}$. So $p \equiv 2 - i \pmod{r}$. Note that it cannot be a valid case for $i = 2$. Then combining equations (3a), (3b) and (3c) with this particular p shows that $(2-i)(k_2 - k_1 + k_3 + 1) \equiv 0 \pmod{r}$. Since r and $2-i$ are coprime ($i \neq 2$), $r|(k_2 - k_1 + k_3 + 1)$, thus $k_2 - k_1 + k_3 + 1 = 0$ or r .
So $(2-i)r^2u_i = -(2-i)r^2$ or 0 . Thus $u_i = -1$ or 0 .

□

APPENDIX A.

The coefficients $(u_j)_{1 \leq j \leq \varphi(n)}$ are the solutions of a system of linear equations whose matricial version is $AU = W$ where $W = [(\Phi_m(\xi^i))^{-1}]_{i \in I_n}$ and

$$A = \begin{bmatrix} 1 & \xi & \dots & \xi^{\varphi(n)-1} \\ \vdots & \vdots & \vdots & \vdots \\ 1 & \xi^i & \dots & (\xi^i)^{\varphi(n)-1} \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}_{i \in I_n} = [(\xi^i)^{j-1}]_{\substack{i \in I_n \\ 1 \leq j \leq \varphi(n)-1}}.$$

A is a Vandermonde matrix so it is invertible since all $(\xi^i)_{i \in I_n}$ are distinct. Thus we can give an explicit resolution of the system : $U = A^{-1}W$. We can find in [15] a paragraph concerning the inversion of a Vandermonde matrix but in our case it does not work that well with A since one row and one column are missing. So we are going to use more direct calculations and write $A^{-1} = (1/\det A)^t \text{com } A$. The Vandermonde determinant $\det A$ is

$$\det A = V((\xi^i)_{i \in I_n}) = \prod_{\substack{i, i' \in I_n \\ i < i'}} (\xi^{i'} - \xi^i).$$

The comatrix has coefficients $[\text{com } A]_{ij} = (-1)^{i+j} V_{ij}$. We call V_{ij} the determinant composed of all the coefficients in A , except the i -th row and the j -th column.

Lemma 2. Let $\sigma_l(a_1, \dots, a_r)$ denote the l -th elementary symmetric function (without the question of the sign).

$$\sigma_l(a_1, \dots, a_r) = \sum_{1 \leq i_1 < \dots < i_l \leq r} a_{i_1} \times \dots \times a_{i_l}.$$

If $A = [a_i^j]_{\substack{1 \leq i \leq r \\ 0 \leq j \leq r-1}}$ then $\det A = V((a_i)_{1 \leq i \leq r})$ (well-known Vandermonde determinant) and its comatrix is given by

$$[com A]_{i_0, j_0} = (-1)^{i_0 + j_0} \sigma_{r-j_0}(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r) V(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r).$$

Proof. All we have to do is write the polynomial $P(X) = \prod_{\substack{1 \leq i \leq r \\ i \neq i_0}} (X - a_i)$ in terms of a Vandermonde determinant and identify the coefficients of the polynomial in X that we obtain.

Let $M(X)$ be simply the block matrix

$$\left[\begin{array}{cccc} 1 & X & \dots & X^{r-1} \\ \hline & A & & \end{array} \right].$$

Now let us expand along the first row the following determinant, consisting in the matrix M without the row $[1 \ a_{i_0} \ \dots \ a_{i_0}^{r-1}]$.

$$|(M_{ij})_{i \neq i_0+1}| = \sum_{k=1}^r (-1)^{k-1} V_{i_0, k} X^{k-1}. \quad (3)$$

Indeed we can write it without loss of generality in a convenient way for $i_0 = r$.

$$\begin{aligned} \begin{vmatrix} 1 & X & \dots & X^{r-1} \\ 1 & a_1 & \dots & a_1^{r-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & a_{r-1} & \dots & a_{r-1}^{r-1} \end{vmatrix} &= \begin{vmatrix} a_1 & \dots & a_1^{r-1} \\ \vdots & \ddots & \vdots \\ a_{r-1} & \dots & a_{r-1}^{r-1} \end{vmatrix} - \begin{vmatrix} 1 & a_1^2 & \dots & a_1^{r-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & a_{r-1}^2 & \dots & a_{r-1}^{r-1} \end{vmatrix} X + \dots \\ &\dots + (-1)^{r-1} \begin{vmatrix} 1 & a_1 & \dots & a_1^{r-2} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & a_{r-1} & \dots & a_{r-1}^{r-2} \end{vmatrix} X^{r-1}. \end{aligned}$$

Besides $|(M_{ij})_{i \neq i_0+1}|$ is nothing but a Vandermonde determinant.

$$\begin{aligned} |(M_{ij})_{i \neq i_0+1}| &= \prod_{\substack{1 \leq i \leq r \\ i \neq i_0}} (a_i - X) \underbrace{\prod_{\substack{1 \leq i < i' \leq r \\ i, i' \neq i_0}} (a_{i'} - a_i)}_{V(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r)} \\ &= (-1)^{r-1} V(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r) \prod_{\substack{1 \leq i \leq r \\ i \neq i_0}} (X - a_i). \end{aligned}$$

But we also know the expansion of this product in terms of symmetric functions (with the convention $\sigma_0 = 1$) :

$$\prod_{\substack{1 \leq i \leq r \\ i \neq i_0}} (X - a_i) = \sum_{k=1}^r (-1)^{r-k} \sigma_{r-k}(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r) X^{k-1}.$$

Identifying this relation with (3), we finally have

$$\forall i_0, j_0, \quad V_{i_0, j_0} = \sigma_{r-j_0}(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r) V(a_1, \dots, \widehat{a_{i_0}}, \dots, a_r),$$

which gives the expected formula since $[\text{com } A]_{i_0, j_0} = (-1)^{i_0+j_0} V_{i_0, j_0}$. $\square$

Corollary 2. *For all $1 \leq i, j \leq \varphi(n)$, the comatrix of A has the following coefficients.*

$$[\text{com } A]_{i,j} = (-1)^{i+j} \sigma_{\varphi(n)-j} \left((\xi^k)_{k \in I_n \setminus \{i\}} \right) V \left((\xi^k)_{k \in I_n \setminus \{i\}} \right).$$

Proof. We just have to use Lemma 2 with $r = \varphi(n)$. The $(a_i)_i$ are $(\xi^i)_{i \in I_n}$. $\square$

Now we can compute the coefficients of U .

Lemma 3.

$$\forall 1 \leq i \leq \varphi(n), \quad u_i = \sum_{j \in I_n} (-1)^{i+j} \sigma_{\varphi(n)-i} \left((\xi^k)_{k \in I_n \setminus \{j\}} \right) \frac{V \left((\xi^k)_{k \in I_n \setminus \{j\}} \right)}{V \left((\xi^k)_{k \in I_n} \right)} \Phi_m(\xi^j)^{-1}.$$

Proof. First the coefficients of A^{-1} are

$$\begin{aligned} [A^{-1}]_{i,j} &= \frac{1}{\det A} [\text{com } A]_{j,i} \\ &= (-1)^{i+j} \sigma_{\varphi(n)-i} \left((\xi^k)_{k \in I_n \setminus \{j\}} \right) \frac{V \left((\xi^k)_{k \in I_n \setminus \{j\}} \right)}{V \left((\xi^k)_{k \in I_n} \right)}. \end{aligned}$$

And now

$$\begin{aligned} u_i &= \sum_{j=1}^{\varphi(n)} [A^{-1}]_{i,j} W_j \\ &= \sum_{j=1}^{\varphi(n)} (-1)^{i+j} \sigma_{\varphi(n)-i} \left((\xi^k)_{k \in I_n \setminus \{j\}} \right) \frac{V \left((\xi^k)_{k \in I_n \setminus \{j\}} \right)}{V \left((\xi^k)_{k \in I_n} \right)} \Phi_m(\xi^j)^{-1}. \end{aligned}$$

$\square$

Remark. The two Vandermonde determinants involved have many terms in common. After simplification there only remains

$$\frac{V \left((\xi^k)_{k \in I_n \setminus \{j\}} \right)}{V \left((\xi^k)_{k \in I_n} \right)} = \pm \frac{\Phi_n(X)}{X - \xi_j} \Big|_{X=\xi_j}.$$

REFERENCES

- [1] T. M. Apostol. Resultants of Cyclotomic Polynomials. *Proceedings of the American Mathematical Society* (1970), Vol. 24, no. 3, pp. 457-462.
- [2] G. Bachman. On the Coefficients of Ternary Cyclotomic Polynomials. *Journal of Number Theory* **100** (2003), pp. 104-116.
- [3] A. S. Bang. Om Ligningen $\phi_n(x) = 0$. *Nyt Tidsskrift for Mathematic* **6** (1895), pp. 6-12.
- [4] P.T. Bateman. Note on the coefficients of the cyclotomic polynomial. *Bulletin of the American Mathematical Society* **55**, 1180-1181.
- [5] M. Beiter. Magnitude of the Coefficients of the Cyclotomic Polynomial F_{pqr} , II. *Duke math. J.* **38** (1971), pp. 591-594.
- [6] D. M. Bloom. On the Coefficients of The Cyclotomic Polynomials. *The American Mathematical Monthly* **75** (1968), pp. 372-377.
- [7] L. Carlitz. The Number of Terms in the Cyclotomic Polynomial F_{pqr} . *The American Mathematical Monthly* **73** (1966), pp. 979-981
- [8] M. van Dijk and D. Woodruff. Asymptotically Optimal Communication for Torus-Based Cryptography. *Crypto'04*, LNCS **3152**, 157-178.
- [9] C. Dunand and R. Lercier. Elliptic Bases and Torus-Based Cryptography. To be published.
- [10] P. Erdős. On the Coefficients of the Cyclotomic Polynomial. *Bulletin of the American Mathematical Society* **52** (1946), pp. 179-184.
- [11] Y. Gallot and P. Moree. Ternary Cyclotomic Polynomials Having a Large Coefficient. arXiv: 0712.2365, Crelle's J., in press.
- [12] T. Y. Lam and K. H. Leung. On the Cyclotomic Polynomial $\Phi_{pq}(X)$, *The American Mathematical Monthly* **103** (1996), pp. 562-564.
- [13] A. Migotti. Zur Theorie der Kreisteilungsgleichung. *S.-B. der Math.-Naturwiss. Classe der Kaiserlichen Akademie der Wissenschaften, Wien* **87** (1883), pp 7-14.
- [14] K. Rubin and A. Silverberg. Torus-Based Cryptography. *Crypto'03*, LNCS **2729**, pp. 349-365.
- [15] J. von zur Gathen and J. Gerhard. *Modern Computer Algebra*. Cambridge University Press (1999).

INSTITUT DE RECHERCHE MATHÉMATIQUE DE RENNES, UNIVERSITÉ DE RENNES 1, CAMPUS DE BEAULIEU,
F-35042 RENNES CEDEX, FRANCE.

E-mail address: `clement.dunand@univ-rennes1.fr`